



# Uncertainty

---

## Chapter 13



# Outline

---

- Uncertainty
- Probability
- Syntax and Semantics
- Inference
- Independence and Bayes' Rule



# Uncertainty

---

Let action  $A_t$  = leave for airport  $t$  minutes before flight  
Will  $A_t$  get me there on time?

Problems:

1. partial observability (road state, other drivers' plans, etc.)
2. noisy sensors (traffic reports)
3. uncertainty in action outcomes (flat tire, etc.)
4. immense complexity of modeling and predicting traffic

Hence a purely logical approach either

1. risks falsehood: " $A_{25}$  will get me there on time", or
2. leads to conclusions that are too weak for decision making:

" $A_{25}$  will get me there on time if there's no accident on the bridge and it doesn't rain and my tires remain intact etc etc."

( $A_{1440}$  might reasonably be said to get me there on time but I'd have to stay overnight in the airport ...)

# Methods for handling uncertainty

- Default or nonmonotonic logic:
  - Assume my car does not have a flat tire
  - Assume  $A_{25}$  works unless contradicted by evidence
- Issues: What assumptions are reasonable? How to handle contradiction?
  
- Rules with fudge factors:
  - $A_{25} \dashv\rightarrow_{0.3}$  get there on time
  - $Sprinkler \dashv\rightarrow_{0.99} WetGrass$
  - $WetGrass \dashv\rightarrow_{0.7} Rain$
- Issues: Problems with combination, e.g., *Sprinkler causes Rain??*
  
- Probability
  - Model agent's degree of belief
  - Given the available evidence,
  - $A_{25}$  will get me there on time with probability 0.04



# Probability

---

Probabilistic assertions **summarize** effects of

- **laziness**: failure to enumerate exceptions, qualifications, etc.
- **ignorance**: lack of relevant facts, initial conditions, etc.

**Subjective** probability:

- Probabilities relate propositions to agent's own state of knowledge

$$\text{e.g., } P(A_{25} \mid \text{no reported accidents}) = 0.06$$

These are **not** assertions about the world

Probabilities of propositions change with new evidence:

$$\text{e.g., } P(A_{25} \mid \text{no reported accidents, 5 a.m.}) = 0.15$$



# Making decisions under uncertainty

---

Suppose I believe the following:

$$P(A_{25} \text{ gets me there on time} \mid \dots) = 0.04$$

$$P(A_{90} \text{ gets me there on time} \mid \dots) = 0.70$$

$$P(A_{120} \text{ gets me there on time} \mid \dots) = 0.95$$

$$P(A_{1440} \text{ gets me there on time} \mid \dots) = 0.9999$$

- Which action to choose?

Depends on my **preferences** for missing flight vs. time spent waiting, etc.

- **Utility theory** is used to represent and infer preferences
- **Decision theory** = probability theory + utility theory



# Syntax

---

- Basic element: **random variable**
- Similar to propositional logic: possible worlds defined by assignment of values to random variables.
- **Boolean** random variables  
e.g., *Cavity* (do I have a cavity?)
- **Discrete** random variables  
e.g., *Weather* is one of  $\langle \textit{sunny}, \textit{rainy}, \textit{cloudy}, \textit{snow} \rangle$
- Domain values must be exhaustive and mutually exclusive
- Elementary proposition constructed by assignment of a value to a random variable: e.g.,  $\textit{Weather} = \textit{sunny}, \textit{Cavity} = \textit{false}$  (abbreviated as  $\neg \textit{cavity}$ )
- Complex propositions formed from elementary propositions and standard logical connectives e.g.,  $\textit{Weather} = \textit{sunny} \vee \textit{Cavity} = \textit{false}$



# Syntax

---


- **Atomic event:** A **complete** specification of the state of the world about which the agent is uncertain  
E.g., if the world consists of only two Boolean variables *Cavity* and *Toothache*, then there are 4 distinct atomic events:
  - $Cavity = false \wedge Toothache = false$
  - $Cavity = false \wedge Toothache = true$
  - $Cavity = true \wedge Toothache = false$
  - $Cavity = true \wedge Toothache = true$
- Atomic events are mutually exclusive and exhaustive



# Axioms of probability

- For any propositions  $A, B$ 
  - $0 \leq P(A) \leq 1$
  - $P(\text{true}) = 1$  and  $P(\text{false}) = 0$
  - $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$

True



# Prior probability

- **Prior or unconditional probabilities** of propositions  
e.g.,  $P(\text{Cavity} = \text{true}) = 0.1$  and  $P(\text{Weather} = \text{sunny}) = 0.72$  correspond to belief prior to arrival of any (new) evidence
- **Probability distribution** gives values for all possible assignments:  
 $P(\text{Weather}) = \langle 0.72, 0.1, 0.08, 0.1 \rangle$  (**normalized**, i.e., sums to 1)
- **Joint probability distribution** for a set of random variables gives the probability of every atomic event on those random variables  
 $P(\text{Weather}, \text{Cavity}) =$  a  $4 \times 2$  matrix of values:

| <i>Weather</i> =      | sunny | rainy | cloudy | snow |
|-----------------------|-------|-------|--------|------|
| <i>Cavity</i> = true  | 0.144 | 0.02  | 0.016  | 0.02 |
| <i>Cavity</i> = false | 0.576 | 0.08  | 0.064  | 0.08 |

- **Every question about a domain can be answered by the joint distribution**



# Conditional probability

---

- Conditional or posterior probabilities  
e.g.,  $P(\text{cavity} \mid \text{toothache}) = 0.8$   
i.e., given that *toothache* is all I know
- Notation for conditional distributions:  
 $P(\text{cavity} \mid \text{toothache}) = 2\text{-element vector of } 2\text{-element vectors}$
- If we know more, e.g., *cavity* is also given, then we have  
 $P(\text{cavity} \mid \text{toothache}, \text{cavity}) = 1$
- New evidence may be irrelevant, allowing simplification, e.g.,  
 $P(\text{cavity} \mid \text{toothache}, \text{sunny}) = P(\text{cavity} \mid \text{toothache}) = 0.8$ 
  - This kind of inference, sanctioned by domain knowledge, is crucial

# Conditional probability

- Definition of conditional probability:  
 $P(a \mid b) = P(a \wedge b) / P(b)$  if  $P(b) > 0$
- **Product rule** gives an alternative formulation:  
 $P(a \wedge b) = P(a \mid b) P(b) = P(b \mid a) P(a)$
- A general version holds for whole distributions, e.g.,  
 $P(\textit{Weather}, \textit{Cavity}) = P(\textit{Weather} \mid \textit{Cavity}) P(\textit{Cavity})$ 
  - (View as a set of  $4 \times 2$  equations, **not** matrix mult.)
- **Chain rule** is derived by successive application of product rule:  
$$\begin{aligned} P(X_1, \dots, X_n) &= P(X_1, \dots, X_{n-1}) P(X_n \mid X_1, \dots, X_{n-1}) \\ &= P(X_1, \dots, X_{n-2}) P(X_{n-1} \mid X_1, \dots, X_{n-2}) P(X_n \mid X_1, \dots, X_{n-1}) \\ &= \dots \\ &= \prod_{i=1}^n P(X_i \mid X_1, \dots, X_{i-1}) \end{aligned}$$

# Inference by enumeration

- Start with the joint probability distribution:

|                      | <i>toothache</i> |                     | $\neg$ <i>toothache</i> |                     |
|----------------------|------------------|---------------------|-------------------------|---------------------|
|                      | <i>catch</i>     | $\neg$ <i>catch</i> | <i>catch</i>            | $\neg$ <i>catch</i> |
| <i>cavity</i>        | <b>.108</b>      | <b>.012</b>         | <b>.072</b>             | <b>.008</b>         |
| $\neg$ <i>cavity</i> | <b>.016</b>      | <b>.064</b>         | <b>.144</b>             | <b>.576</b>         |

- For any proposition  $\phi$ , sum the atomic events where it is true:  $P(\phi) = \sum_{\omega:\omega \models \phi} P(\omega)$

# Inference by enumeration

- Start with the joint probability distribution:

|                      | <i>toothache</i> |                     | $\neg$ <i>toothache</i> |                     |
|----------------------|------------------|---------------------|-------------------------|---------------------|
|                      | <i>catch</i>     | $\neg$ <i>catch</i> | <i>catch</i>            | $\neg$ <i>catch</i> |
| <i>cavity</i>        | <b>.108</b>      | <b>.012</b>         | <b>.072</b>             | <b>.008</b>         |
| $\neg$ <i>cavity</i> | <b>.016</b>      | <b>.064</b>         | <b>.144</b>             | <b>.576</b>         |

- For any proposition  $\varphi$ , sum the atomic events where it is true:  $P(\varphi) = \sum_{\omega:\omega \models \varphi} P(\omega)$
- $P(\textit{toothache}) = 0.108 + 0.012 + 0.016 + 0.064 = 0.2$

# Inference by enumeration

- Start with the joint probability distribution:

|                      | <i>toothache</i> |                     | $\neg$ <i>toothache</i> |                     |
|----------------------|------------------|---------------------|-------------------------|---------------------|
|                      | <i>catch</i>     | $\neg$ <i>catch</i> | <i>catch</i>            | $\neg$ <i>catch</i> |
| <i>cavity</i>        | <b>.108</b>      | <b>.012</b>         | <b>.072</b>             | <b>.008</b>         |
| $\neg$ <i>cavity</i> | <b>.016</b>      | <b>.064</b>         | <b>.144</b>             | <b>.576</b>         |

- For any proposition  $\phi$ , sum the atomic events where it is true:  $P(\phi) = \sum_{\omega: \omega \models \phi} P(\omega)$

- $P(\textit{toothache}) = 0.108 + 0.012 + 0.016 + 0.064 = 0.2$

# Inference by enumeration

- Start with the joint probability distribution:

|                      | <i>toothache</i> |                     | $\neg$ <i>toothache</i> |                     |
|----------------------|------------------|---------------------|-------------------------|---------------------|
|                      | <i>catch</i>     | $\neg$ <i>catch</i> | <i>catch</i>            | $\neg$ <i>catch</i> |
| <i>cavity</i>        | .108             | .012                | .072                    | .008                |
| $\neg$ <i>cavity</i> | .016             | .064                | .144                    | .576                |

- Can also compute conditional probabilities:

$$\begin{aligned} P(\neg \text{cavity} \mid \text{toothache}) &= \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} \\ &= \frac{0.016 + 0.064}{0.108 + 0.012 + 0.016 + 0.064} \\ &= 0.4 \end{aligned}$$



# Normalization

|                      | <i>toothache</i> |                     | $\neg$ <i>toothache</i> |                     |
|----------------------|------------------|---------------------|-------------------------|---------------------|
|                      | <i>catch</i>     | $\neg$ <i>catch</i> | <i>catch</i>            | $\neg$ <i>catch</i> |
| <i>cavity</i>        | <b>.108</b>      | <b>.012</b>         | .072                    | .008                |
| $\neg$ <i>cavity</i> | <b>.016</b>      | <b>.064</b>         | .144                    | .576                |

- Denominator can be viewed as a **normalization constant**  $\alpha$

$$\begin{aligned}
 P(\textit{Cavity} \mid \textit{toothache}) &= \alpha \cdot P(\textit{Cavity}, \\
 &\quad \textit{toothache}) \\
 &= \alpha \cdot [P(\textit{Cavity}, \textit{toothache}, \textit{catch}) + P(\textit{Cavity}, \\
 &\quad \textit{toothache}, \neg \textit{catch})]
 \end{aligned}$$



# Inference by enumeration, contd.

---

Typically, we are interested in the posterior joint distribution of the **query variables**  $Y$  given specific values  $e$  for the **evidence variables**  $E$

Let the **hidden variables** be  $H = X - Y - E$


Then the required summation of joint entries is done by summing out the hidden variables:

$$P(Y \mid E = e) = \alpha P(Y, E = e) = \alpha \sum_{\mathbf{h}} P(Y, E = e, H = \mathbf{h})$$

- The terms in the summation are joint entries because  $Y$ ,  $E$  and  $H$  together exhaust the set of random variables
- Obvious problems:
  1. Worst-case time complexity  $O(d^n)$  where  $d$  is the largest arity
  2. Space complexity  $O(d^n)$  to store the joint distribution
  3. How to find the numbers for  $O(d^n)$  entries?

# Independence

- $A$  and  $B$  are independent iff  
 $P(A/B) = P(A)$  or  $P(B/A) = P(B)$  or  $P(A, B) = P(A) P(B)$



$$P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}, \textit{Weather}) \\ = P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}) P(\textit{Weather})$$

- 32 entries reduced to 12 (8+4);  
for  $n$  independent biased coins,  $O(2^n) \rightarrow O(n)$
- Absolute independence powerful but rare
- Dentistry is a large field with hundreds of variables, none of which are independent. What to do?



# Conditional independence

---

- $P(\textit{Toothache}, \textit{Cavity}, \textit{Catch})$  has  $2^3 - 1 = 7$  independent entries
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:  
(1)  $P(\textit{catch} \mid \textit{toothache}, \textit{cavity}) = P(\textit{catch} \mid \textit{cavity})$
- The same independence holds if I haven't got a cavity:  
(2)  $P(\textit{catch} \mid \textit{toothache}, \neg\textit{cavity}) = P(\textit{catch} \mid \neg\textit{cavity})$
- *Catch* is **conditionally independent** of *Toothache* given *Cavity*:  
 $P(\textit{Catch} \mid \textit{Toothache}, \textit{Cavity}) = P(\textit{Catch} \mid \textit{Cavity})$
- Equivalent statements:  
 $P(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) = P(\textit{Toothache} \mid \textit{Cavity})$   
 $P(\textit{Toothache}, \textit{Catch} \mid \textit{Cavity}) = P(\textit{Toothache} \mid \textit{Cavity}) P(\textit{Catch} \mid \textit{Cavity})$

# Conditional independence contd.

- Write out full joint distribution using chain rule:

$$\begin{aligned} P(\textit{Toothache}, \textit{Catch}, \textit{Cavity}) &= P(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) P(\textit{Catch}, \textit{Cavity}) \\ &= P(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) P(\textit{Catch} \mid \textit{Cavity}) P(\textit{Cavity}) \\ &= P(\textit{Toothache} \mid \textit{Cavity}) P(\textit{Catch} \mid \textit{Cavity}) P(\textit{Cavity}) \end{aligned}$$

I.e.,  $2 + 2 + 1 = 5$  independent numbers

- In most cases, the use of conditional independence reduces the size of the representation of the joint distribution from exponential in  $n$  to linear in  $n$ .
- Conditional independence is our most basic and robust form of knowledge about uncertain environments.

# Bayes' Rule

- Product rule  $P(a \wedge b) = P(a | b) P(b) = P(b | a) P(a)$   
 $\Rightarrow$  Bayes' rule:  $P(a | b) = P(b | a) P(a) / P(b)$
- or in distribution form  
$$P(Y|X) = P(X|Y) P(Y) / P(X) = \alpha P(X|Y) P(Y)$$
- Useful for assessing **diagnostic** probability from **causal** probability:
  - $P(\text{Cause}|\text{Effect}) = P(\text{Effect}|\text{Cause}) P(\text{Cause}) / P(\text{Effect})$
  - E.g., let  $M$  be meningitis,  $S$  be stiff neck:  
 $P(m|s) = P(s|m) P(m) / P(s) = 0.5 \times 0.0002 / 0.05 = 0.0002$
  - Note: posterior probability of meningitis still very small!

# Bayes' Rule and conditional independence

$$P(\text{Cavity} \mid \text{toothache} \wedge \text{catch})$$

$$= \alpha \cdot P(\text{toothache} \wedge \text{catch} \mid \text{Cavity}) P(\text{Cavity})$$

$$= \alpha \cdot P(\text{toothache} \mid \text{Cavity}) P(\text{catch} \mid \text{Cavity}) P(\text{Cavity})$$

- This is an example of a **naïve Bayes** model:

$$P(\text{Cause}, \text{Effect}_1, \dots, \text{Effect}_n) = P(\text{Cause}) \prod_i P(\text{Effect}_i \mid \text{Cause})$$



- Total number of parameters is **linear** in  $n$



# Summary

---

- Probability is a rigorous formalism for uncertain knowledge
- **Joint probability distribution** specifies probability of every **atomic event**
- Queries can be answered by summing over atomic events
- For nontrivial domains, we must find a way to reduce the joint size
- **Independence** and **conditional independence** provide the tools